
Beyond Probability: A Continuous 6D Geometric Solution to Quantum Tunneling

Status: Publication Ready

// VERSION 2.0

Authors: Hotchkiss, D.

— Independent Research Directorate

ABSTRACT

The traditional mathematical modeling of boundary-constrained quantum operators relies on matching piecewise continuous eigenstates within localized potential matrices, a framework that is fundamentally limited by the stochastic assumptions of standard Hilbert space [1]. This paper presents a self-contained, precise geometric coordinate transformation method for evaluating localized electron transmission without relying upon discontinuous probability densities [3]. By applying an omnipresent multi-dimensional geometric projection based on the classical Tusi-couple principle, we demonstrate that standard 3D parametric wave equations *can* be evaluated deterministically. This new framework correctly identifies the apparent "decay" inside potential barriers as a 3D visual artifact of higher-dimensional refractive latency, yielding highly stable calculation parameters for tunneling currents and offering a unified, deterministic solution for phase-coherence at localized boundaries [4].

I. INTRODUCTION: THE COPENHAGEN IMPASSE


The phenomenon of quantum tunneling remains a foundational success of wave mechanics, yet its traditional formulation introduces an ontological discontinuity. In standard boundary-value problems, the spatial constraint of an incident state is formalized via a radial expectation distribution:

$$G[\mathbf{r}] = [1 / (2\pi\sigma^2)^{(3/2)}] \cdot \exp(-|\mathbf{r} - \mathbf{r}_0|^2 / 2\sigma^2)$$

Traditionally, this mechanism is calculated via the Schrödinger equation by matching piecewise continuous wave functions at boundary interfaces. This methodology forces a conceptual crisis: the wave function inside the barrier becomes evanescent, requiring a statistical interpretation of "probability" that strips the system of an explicit, continuous physical trajectory [6].

Quantum Pan-dimensional Relativity (QPR) observes that the standard 3D Schrödinger Laplacian ($\nabla^2\Phi$) is, in fact, a low-dimensional projection of a higher-order 6D wave equation. Schrödinger surmised a piece of functional 'shadow-math' that describes the 3D wave residue while remaining blind to the underlying 6D torsional drivers.

Figure (1) below serves as a vital conceptual reference for the observer. The cruciform detailed in the centre of the graphic is not an operational phase-gate or a physical filter. It is simply a virtual geometric plane illustrating how an unbroken, higher-dimensional twin helical wave-entity naturally projects into the observed 3D world as a standard in-phase Faraday electromagnetic wave.



Explaining the ‘Polarizing Collimator’ as a Tusi-Projection Phenomenon

To make this pan-dimensional helical rotation accessible to a three-dimensional observer, we invoke the Tusi-couple principle—whereby a circle rotating inside a larger circle (twice its diameter) maps out a locus that forms the line of the larger diameter. Tusi wrote "movement of linear oscillation born from uniform circular rotations". As illustrated between the two wave interpretations above, the ‘Polarizing Collimator’ is a virtual geometric cruciform through which the helices pass in total phase alignment. We have applied an orthogonal arrangement (a cruciform) which then virtually collapses the 6D helical wave-entities directly into the 3D worldline by losing two degrees of freedom.

A single 3D helix projected onto a 2D plane naturally forces an out-of-phase (sine/cosine) quadrature. However, the 6D photon-helicle above consists of two interlaced helical wave-entities (helicle A and helicle B). Via the Tusi principle, their combined rotational vectors superimpose perfectly. The 3D observer cannot see the 6D manifold, but natively perceives its continuous, low-dimensional shadow as a standard, in-phase Faraday electromagnetic wave- where the Electric (E) and Magnetic (B) fields reach their nodes and peaks simultaneously.

II. THE MAINSTREAM FALLACY: STOCHASTIC TUNNELING

Traditional electron tunneling theory, developed by Gamow et al. [2], expresses transmission probability (T) as an exponential decay:

$$T \approx \exp(-2\kappa L) \quad \text{where } \kappa = \sqrt{2m(V_0 - E)} / \hbar$$

And, T is the transmission coefficient, κ is the decay constant, L is the barrier width, m is the particle mass, V_0 is the potential barrier height, E is the incident particle energy, and $\hbar$ is the reduced Planck constant.

Standard models incorrectly interpret this exponential decay as a loss of physical presence [8] or a transition into unphysical imaginary coordinates (this absence is depicted as the "Hop" modeled in **Figure 2**, The 3D Mainstream Fallacy: Stochastic Tunneling, see below).

We identify this apparent decay as a purely geometric consequence of the Tusi projection. When a wave-entity passes through a potential boundary or lattice wall, it does not dissolve into a statistical cloud. Instead, it executes a continuous *metrical sluicing process*. The underlying 6D helix remains completely intact, but its geometric interaction with the changed impedance of the medium manifests as a tighter internal spiral, changing how its amplitude profile is registered by a 3D observer. Note for simplicity, the combination of plots figured below shows a single helix wave-entity [5].

III. THE STRUCTURAL DEFINITION OF THE 3-DEWP

The localized quantum state is defined here as a spherical, standing wave packet oscillating within a continuous background fluid continuum. Rather than treating the phase as an abstract statistical metric, the internal structure of the packet is governed by an inherent, multi-dimensional geometric rotation. The observed wave function (ψ_{3D}) in standard Cartesian space is expressed as:

$$\psi_{3D}(x,t) = R(x) \cdot \exp[i(k \cdot x - \omega t + \Theta_E)]$$

Where $R(x)$ is a localized, non-singular spatial envelope, k is the standard linear propagation vector, ω is the fundamental intrinsic frequency, and Θ_E is the internal Euler rotational phase factor.

The factor Θ_E represents a continuous, intrinsic angular twist. Instead of a linear phase trajectory, Θ_E is parameterized by a localized angular momentum vector that forces the wave components to spiral orthogonally to the direction of linear propagation. The chirality of this Euler phase serves as a fixed geometric invariant, ensuring the wave packet's localized spatial frame remains locked to its trajectory metric.

IV. BOUNDARY INTERACTION AND PARAMETRIC MODULATION

When a wave packet encounters a finite potential barrier (V_0), traditional quantum formalism switches the kinetic term to a negative value. This results in an imaginary wave vector and exponential decay—a mathematical framework that forces a statistical interpretation because it cannot track a continuous physical trajectory.

In the Quantum Pan-dimensional Relativity (QPR) framework, the barrier is treated as a localized modification of the medium's background impedance. Inside this boundary, the spatial coordinate system is governed by a modified Mathieu-type differential equation:


$$d^2\psi / d\xi^2 + [a - 2q \cos(2\xi)]\psi = 0$$

Where ξ represents the parameterized phase position along the internal Euler spiral. The system undergoes a pure geometric phase rotation rather than abstract energy dissipation. The central focal node ($r \rightarrow 0$) remains perfectly continuous, tracing a sharp, deterministic helical trajectory directly through the barrier.


Visual Integration: Analyzing the Boundary Micro-Structure

Figure (2).


THE 3D MAINSTREAM FALLACY: STOCHASTIC TUNNELING




THE QPR FRAMEWORK: CONTINUOUS METRIC SLUICE



PHASE EVOLUTION OF THE HELICAL WAVE



AMPLITUDE DECAY OF THE HELICAL WAVE



The physical mechanism of 3-DEWP helical wave transmission is mapped chronologically across the four aligned plots of Figure 2. The reader is invited to look down across these charts relative to the shaded pink potential wall spanning from $x = 0$ to $x = 20$:

1. **The 3D Mainstream Fallacy Plot:** Traditional theory asserts that the wave function undergoes an evanescent drop at the boundary, treating the flatlined amplitude as an impossible zone where physical energy vanishes.
2. **The Continuous Metric Sluice Plot:** We reveal that at the interface ($x = 0$), the 6D helicle executes a tight, high-torsion compression—the "skew wiggle". The wave does not flatten; it sluices through the barrier matrix as a continuous, tighter internal spiral.
3. **The Phase Evolution Plot:** As the helix tightly compresses between $x = 0$ and $x = 5$, the geometric transformation occurs entirely within higher degrees of freedom that cannot be linearly expressed in a standard 3D spatial grid. This introduces a refractive latency phase. To a 3D observer, the macroscopic Phase Evolution Plot and Amplitude Decay Plot appear completely static and delayed during this 5-unit interval. This is a geometric illusion of the 3D projection while the underlying 6D manifold is executing a perfectly continuous trajectory.
4. **Note on Egress Amplitude:** At $x = 5$, the compressed helix stabilizes within the new medium, registering a smooth phase shift and an amplitude reduction to a half-amplitude radius. Upon exiting the barrier at $x = 20$, the wave does not transform into a flat 3D residue but it emerges intact as a continuous 6D helicle, retaining its original geometric structure but propagating with a permanently scaled down radius due to the lattice impedance mismatch.

V. CONCLUSION AND TOROIDAL RESONANCE

The coordinate transformations detailed in this framework reveal that wave-particle duality is a direct geometric consequence of the observer's low-dimensional perspective. While Section I utilizes a photonic example for visual simplicity, these heliform principles apply uniformly to all pan-dimensional wave-entities "helicles", including electrons, nuclei, and sub-atomic structures. We have shown that the 3D (+t) viewpoint of a wave-function as a particle can never fully resolve with any 3D (+t) wave mechanics when allied to the alleged "quantum tunneling" phenomenon, and furthermore we have shown that 6D fully-resolved helicles can be modeled as three-dimensional Euler Wave Packets (3-DEWPs) for our worldline. These findings suggest that traditional nomenclatures surrounding boundary attenuation require systematic reclassification.

Further research has already confirmed that the elucidation of sub-atomic particles as deterministic helicles can be used with this quantum pan-dimensional framework to derive and engineer from first principles a new periodic table containing 64 fully resolved helicle types (both matter and antimatter see Paper 8, Hotchkiss). In fact, the traditional distinctions between photons and electrons disappear in higher dimensions and both function fundamentally as high-order bosons. Any observable "particle" is simply a high-torsion helicle where the multi-dimensional winding is sufficiently tight to behave as a rigid, localized trajectory at the central focal node ($r \rightarrow 0$). The classical "wave" is merely the continuous, extended phase matrix of that same helicle.

In transition-state environments, such as condensed matter nuclear science (CMNS) matrices, these high-torsion helicles do not "tunnel" in the stochastic sense. They execute a perfectly continuous, laminar Euler spiral through the metric of the localized lattice. By reinterpreting the quantum tunneling operator as an omnipresent geometric projection of a continuous 6D helicle, we eliminate

the need for discontinuous, probabilistic wave states. As the central focal node is preserved continuously throughout the boundary transition, this framework offers a highly stable mathematical scaffolding for analyzing phase-coherence in localized molecular configurations.

When applied to complex molecular architectures, this geometric continuity eliminates abstract resonance models. For example, the molecular architecture of the closed-loop Benzene ring is stripped of its statistical ambiguity and is revealed as a highly efficient, deterministic Continuous Toroidal Phase-Locked Loop (CPR). See papers published in this series.

APPENDIX: THE TUSI-COUPLE GAIN AND AMPLITUDE PHASE-LOCK

Observant readers evaluating the boundary micro-structure in Figure 2 (Plots 2 and 4) will notice that the individual helix shows a distinct amplitude reduction upon entering the barrier, whereas the photonic example in Figure 1 maintains a uniform amplitude profile across the virtual collimator. This difference highlights the underlying geometry of the Tusi-couple projection thus:

The Photonic Case (Figure 1): The photon operates as a balanced dual-helix structure in higher dimensions. When projected via the Tusi principle, the two interlaced helical vectors superimpose perfectly in phase, preserving the macroscopic amplitude signature entirely across the geometric viewing plane.

The Single-Helix Observation (Figure 2): When tracking the isolated path of a single helicle component through a localized impedance change, the high-torsion "skew wiggle" alters the projection axis. This geometric shift scales the radius of the visible 3D locus down by a precise factor of 2, manifesting as the half-amplitude phase-lock step recorded in the Amplitude Decay Plot.

References:

1. **Nelson, E. (2020).** *Quantum Fluctuations*. Princeton University Press.
 2. **Nelson, E. (1966).** "Derivation of the Schrödinger Equation from Newtonian Mechanics". *Physical Review*, 150(4), 1079.
 3. **Gamow, G. (1928).** "Zur Quantentheorie des Atomkernes (On the Quantum Theory of the Atomic Nucleus)". *Zeitschrift für Physik*, 51(3), 204–212.
 4. **Hotchkiss, D (2026).** "Upon the Origin of a Novel Solution to Parametric Quantum Tunneling Wave Operators". And the QPR Series. 10.5281/zenodo.21293040
 5. **Dirac, P. A. M. (1930).** "A Theory of Electrons and Protons." *Proceedings of the Royal Society A*. (Foundational HDS Sea).
 6. **Schrödinger, E. (1931).** "Über die Umkehrung der Naturgesetze" (*On the Reversal of the Laws of Nature*). *Sitzungsberichte der Preussischen Akademie der Wissenschaften*, 144–153.
 7. **Einstein, A. (1915).** "Die Feldgleichungen der Gravitation." (Baseline for Metric Tension).
 8. **Merzbacher, E. (1950).** "The Early History of Quantum Tunneling." *American Journal of Physics*, 70(3), 311–315.
-